arXiv:hep-th/0610040v2 3 Feb 2008

CENTRE DE PHYSIQUE THÉORIQUE [1]
CNRS–Luminy, Case 907
13288 Marseille Cedex 9
FRANCE

On a Classification of Irreducible Almost-Commutative Geometries IV

Jan–Hendrik Jureit [2], Christoph A. Stephan [3]

Abstract

In this paper we will classify the finite spectral triples with *KO*-dimension six, following the classification found in [1, 2, 3, 4], with up to four summands in the matrix algebra. Again, heavy use is made of Krajewski diagrams [5]. This work has been inspired by the recent paper by Alain Connes [6] and John Barrett [7].

In the classification we find that the standard model of particle physics in its minimal version fits the axioms of noncommutative geometry in the case of *KO*-dimension six. By minimal version it is meant that at least one neutrino has to be massless and mass-terms mixing particles and antiparticles are prohibited.

CPT-P75-2006
PACS-92: 11.15 Gauge field theories
MSC-91: 81T13 Yang-Mills and other gauge theories

---

[1] Unité Mixte de Recherche (UMR 6207) du CNRS et des Universités Aix–Marseille 1 et 2 et Sud Toulon–Var, Laboratoire affilié à la FRUMAM (FR 2291)

[2] and Université de Provence and Universität Kiel,
jureit@cpt.univ-mrs.fr

[3] and Université de Provence and Universität Kiel,
stephan@cpt.univ-mrs.fr

1 Introduction

Recently Alain Connes [6] and John Barrett [7] proposed to change the KO-dimension for the finite part of almost-commutative spectral triples from zero to six. Based on this assumption they constructed a version of the standard model of particle physics which allowed for right-handed massive neutrinos in every generation of fermions and a Majorana-mass resulting in the see-saw-mechanism. Furthermore the long standing problem of fermion-doubling could be cured. In the case of KO-dimension six it is possible to directly project out the superfluous degrees of freedom, as is shown in detail in [7].

The price which had to be paid is that not all the axioms of noncommutative geometry [8, 9] are satisfied by this model, notably the orientability axiom which fails on the Lepton-sector [10]. Also the Poincaré duality needs to be modified in the sense that the leptonic sector and the quark sector provide two separate generators of K-homology. Each of these sectors fulfils the Poincaré duality [10].

In this paper we will assume that all the axioms of noncommutative geometry hold and classify the corresponding finite spectral triples following [1, 2, 3, 4]. This classification is based on the classification of finite spectral triples of Mario Paschke, Andrzej Sitarz and Thomas Krajewski [11, 5]. The main tool used to find the possible spectral triples are Krajewski diagrams [5] which have already been used in [1, 2, 3, 4]. Passing from KO-dimension zero to KO-dimension six implies a few changes in the definition of a real, finite spectral triple. It will be shown in detail that every real, finite spectral triple in KO-dimension six still allows to split the Hilbert space into left- and right-handed particle and antiparticle subspaces, although this choice may be ambiguous and it does not in principle forbid lepto-quark-like mass terms.

Imposing the axioms of noncommutative geometry will lead us to seven possible minimal Krajewski diagrams, two of which contain the first family of the standard model of particle physics in its minimal version. Thus, if one requires all the axioms to hold, one has to abandon Majorana-masses for right-handed neutrinos and at least one neutrino has to remain massless. This should be compared with the case in KO-dimension zero [1, 2, 3, 4], where 66 Krajewski diagrams appeared, all corresponding to noncommutative geometries which obey to the axioms. It is quite remarkable that a reduction in the input, i.e. no S^0-real structure has to be assumed, does lead to a reduction in the number of possible geometries.

A version of the standard model with four summands in the matrix algebra, with right-handed neutrinos and Majorana-masses has been treated in detail in [14]. This necessitated a modification of the axioms of noncommutative geometry, notably the orientability axiom.

2 Basic Definitions

In this classification we are interested in real, finite spectral triples with KO-dimension six and metric dimension zero, [12, 9]. The metric dimension being zero follows from the requirement of finiteness since this implies that the internal Dirac operator has only a finite number of eigenvalues. Note that no S^0-real structure is imposed. We will show that the axioms still allow to split the Hilbert space into left-handed and right-handed particle and anti-particle subspaces and that the S^0-real structure, which is needed in KO-dimension zero to perform this split, follows naturally from the axioms. It follows also that the Dirac operator must not contain mass terms which connect particles to antiparticles.

Definition 2.1. A real, finite spectral triple of KO-dimension six is given by $(\mathcal{A}, \mathcal{H}, \mathcal{D}, J, \chi)$ with a finite dimensional real algebra $\mathcal{A}$, a faithful representation ρ of $\mathcal{A}$ on a finite dimensional complex Hilbert space $\mathcal{H}$. Three additional operators are defined on $\mathcal{H}$: the Dirac operator $\mathcal{D}$ is selfadjoint, the real structure J is antiunitary, and the chirality χ which is an unitary involution. These operators satisfy:

- $$J^2 = \chi^2 = 1, \quad [J, \mathcal{D}] = \{J, \chi\} = 0, \quad \mathcal{D}\chi = -\chi\mathcal{D}, \quad [\chi, \rho(a)] = 0,$$
  $$[\rho(a), J\rho(b)J^{-1}] = [[\mathcal{D}, \rho(a)], J\rho(b)J^{-1}] = 0, \forall a, b \in \mathcal{A}. \qquad (2.1)$$

Note that in KO-dimension six the commutator $[J, \chi] = 0$ from KO-dimension zero becomes an anti-commutator [6, 7].

- The chirality can be written as a finite sum $\chi = \sum_i \rho(a_i) J\rho(b_i) J^{-1}$. This condition is called orientability. The finite sum is a zero dimensional Hochschild cycle.

- The intersection form $\cap_{ij} := \mathrm{tr}(\chi\, \rho(p_i) J\rho(p_j) J^{-1})$ is non-degenerate, $\det \cap \neq 0$. The p_i are minimal rank projections in $\mathcal{A}$. This condition is called *Poincaré duality.*

The algebra is a finite sum of N simple algebras , $\mathcal{A} = \oplus_{i=1}^N \mathcal{A}_i$ with $\mathcal{A}_i = M_{n_i}(\mathbb{K}_i)$, and $\mathbb{K}_i = \mathbb{R}, \mathbb{C}, \mathbb{H}$ where $\mathbb{H}$ denotes the quaternions.

We will now give a derivation of the substructure of the operators and the Hilbert space constituting the spectral triple. It will turn out that the vanishing anti-commutator $\{J, \chi\} = 0$ and the axiom of orientability replace the S^0-real structure of the case with KO-dimension zero.

The chirality, the real structure and the Hilbert space: Since the chirality is a unitary operator with $\chi^2 = 1$ we can build the projectors $(1 \mp \chi)/2$ which allow to decompose the Hilbert space

$$\mathcal{H} = \mathcal{H}^- \oplus \mathcal{H}^+ . \qquad (2.2)$$

The first component corresponds in physics to left-handed particles and to charge conjugate right-handed particles, $\chi = -1$, the second component corresponds to right-handed particles and the charge conjugate of left-handed particles, $\chi = +1$. The real structure anti-commutes with the chirality, $\{J, \chi\} = 0$, therefore it maps the subspace $\mathcal{H}^-$ to the subspace $\mathcal{H}^+$ and vice versa. And since $J^2 = 1$ we have $\dim\mathcal{H}^- = \dim\mathcal{H}^+ = n$. As a convention we will take the basis of $\mathcal{H}$ in which the chirality is a diagonal matrix with eigenvalues ∓ 1

$$\chi = (-1_n) \oplus 1_n \qquad (2.3)$$

and in which the real structure takes the standard form

$$J = \begin{pmatrix} 0 & 1_n \\ 1_n & 0 \end{pmatrix} \circ \text{complex conjugation.} \tag{2.4}$$

The representation: The chirality and the real structure automatically require the representation ρ of the algebra $\mathcal{A}$ to be block-diagonal on the subspaces $\mathcal{H}^-$ and $\mathcal{H}^+$

$$\rho(a) = \rho^-(a) \oplus \rho^+(a), \quad a \in \mathcal{A}. \tag{2.5}$$

Concerning the algebra $\mathcal{A} = \oplus_{i=1}^N \mathcal{A}_i$, we restrict ourselves to the easy case, $\mathbb{K} = \mathbb{R}, \mathbb{H}$ in all components $\mathcal{A}_i = M_{n_i}(\mathbb{K}_i)$ of the algebra. The sub-algebras $M_n(\mathbb{R})$ and $M_n(\mathbb{H})$ only have one irreducible representation, the fundamental one on $\mathbb{C}^{(n)}$, where $(n) = n$ for $\mathbb{K} = \mathbb{R}$ and $(n) = 2n$ for $\mathbb{K} = \mathbb{H}$. All of the following arguments also hold for $\mathbb{K} = \mathbb{C}$, but the notation becomes more opaque since the complex conjugate of the fundamental representation has to be taken into account. We notice that the axiom $[\rho(a), J\rho(b)J^{-1}] = 0$ for all $a, b \in \mathcal{A}$ requires $\rho(a)$ to be of the form

$$\rho(a) = \left(\oplus_{i,j=1}^N \rho^-(a_i, a_j)\right) \oplus \left(\oplus_{i,j=1}^N \rho^+(a_i, a_j)\right) \tag{2.6}$$

where ρ^- and ρ^+ can be either of the form

$$\rho^-(a_i, a_j) := a_i \otimes 1_{m_{ij}} \otimes 1_{n_j} \quad \rho^+(a_i, a_j) := 1_{n_i} \otimes 1_{m_{ij}} \otimes a_j \tag{2.7}$$

or of the form

$$\rho^-(a_i, a_j) := 1_{n_i} \otimes 1_{m_{ij}} \otimes a_j \quad \rho^+(a_i, a_j) := a_i \otimes 1_{m_{ij}} \otimes 1_{n_j}. \tag{2.8}$$

The multiplicities m_{ij} are non-negative integers and we denote by 1_n the $n \times n$ identity matrix and set by convention $1_0 := 0$. Algebra elements a_i are taken to be from he ith summand $M_{n_i}(\mathbb{K}_i)$ of the algebra $\mathcal{A} = \oplus_{i=1}^N M_{n_i}(\mathbb{K}_i)$.

The real structure J permutes the two main summands ρ^- and ρ^+ of ρ (2.6) and complex conjugates them. The chirality reads explicitly

$$\chi \;=\; (\oplus_{i,j=1}^N 1_{n_i} \otimes (-1) 1_{m_{ij}} \otimes 1_{n_j}) \;\oplus\; (\oplus_{i,j=1}^N 1_{n_i} \otimes 1_{m_{ij}} \otimes 1_{n_j}). \tag{2.9}$$

To simplify the following calculations we will also assume $m_{ij} = 1$ in all sub-representations.

The orientability axiom allows to put further restrictions on the sub-representations $\rho(a_i, a_j) = \rho^-(a_i, a_j) \oplus \rho^+(a_i, a_j)$. Let χ_{ij} and J_{ij} be the chirality and the real structure restricted to the subspace $\mathcal{H}_{ij} = \mathcal{H}_{ij}^- \oplus \mathcal{H}_{ij}^+$ corresponding to $\rho(a_i, a_j)$. Then the orientability axiom requires that there exist $a_{i,r}, b_{i,r} \in \mathcal{A}_i$ and $a_{j,r}, b_{j,r} \in \mathcal{A}_j$ such that

$$\sum_r \rho(a_{i,r}, a_{j,r}) J_{ij} \rho(b_{i,r}, b_{j,r}) J_{ij}^{-1} \;=\; \chi_{ij}. \tag{2.10}$$

Without loss of generality we can choose the sub-representation to be of the form 2.7 and find in a straightforward calculation

$$\sum_r \rho(a_{i,r}, a_{j,r}) J_{ij} \rho(b_{i,r}, b_{j,r}) J_{ij}^{-1} \;= \sum_r (a_{i,r} \otimes b_{j,r}) \oplus (b_{i,r} \otimes a_{j,r}) \tag{2.11}$$

which is required to be equal to $\chi_{ij} = (-1)1_{(n_i+n_j)} \oplus 1_{(n_i+n_j)}$. It follows immediately that $i \neq j$. The same argument holds for the complex case, $\mathbb{K} = \mathbb{C}$.

Note that this argument forbids Majorana masses for right-handed neutrinos if the orientability axiom is used in its classical form [14].

The Dirac operator: The Dirac operator anti-commutes with the chirality, $\{\mathcal{D}, \chi\} = 0$, and maps therefore the sub-space $\mathcal{H}^-$ to $\mathcal{H}^+$ and vice versa. Taking into account the self-adjointness of $\mathcal{D}$ leads to the following general form

$$\mathcal{D} = \begin{pmatrix} 0 & \Delta \\ \Delta^* & 0 \end{pmatrix}, \quad \text{with } \Delta : \mathcal{H}^- \to \mathcal{H}^+. \tag{2.12}$$

Furthermore $\mathcal{D}$ commutes with the real structure J,

$$[\mathcal{D}, J] = \begin{pmatrix} \overline{\Delta}^* - \Delta & 0 \\ 0 & \overline{\Delta} - \Delta^* \end{pmatrix} = 0 \tag{2.13}$$

and therefore $\Delta^t = \Delta$. And so the Dirac operator reads:

$$\mathcal{D} = \begin{pmatrix} 0 & \Delta \\ \overline{\Delta} & 0 \end{pmatrix} \quad \text{with } \Delta^t = \Delta. \tag{2.14}$$

This form of the Dirac operator seems to deviate from the standard one used in the literature, but this is only due to the choice of basis which is obtained by ordering the particle and anti-particle multiplets with respect to their chirality.

The first order axiom: We will now employ the first order axiom,

$$[[\mathcal{D}, \rho(a)], J\rho(b)J^{-1}] = 0, \forall a, b \in \mathcal{A}, \tag{2.15}$$

to put further restrictions on the fine structure of the Dirac operator and the representation. Using the equation 2.14 for the Dirac operator and the representation 2.5 one finds with

$$J\rho(a)J^{-1} = \overline{\rho}^-(a) \oplus \overline{\rho}^+(a) \tag{2.16}$$

after a simple calculation the following two equations for the first order axiom

$$\Delta\, \rho^-(a)\, \overline{\rho}^+(b) - \rho^+(a)\, \Delta\, \rho^+(b) - \overline{\rho}^-(b)\, \Delta\, \rho^-(a) + \overline{\rho}^-(b)\, \rho^+(a)\, \Delta = 0 \tag{2.17}$$

$$\overline{\Delta}\, \rho^+(a)\, \overline{\rho}^-(b) - \rho^-(a)\, \overline{\Delta}\, \rho^-(b) - \overline{\rho}^+(b)\, \overline{\Delta}\, \rho^+(a) + \overline{\rho}^+(b)\, \rho^-(a)\, \overline{\Delta} = 0, \tag{2.18}$$

for all $a, b \in \mathcal{A}$. One notices immediately that these two equations are equivalent and we will therefore use from now on the first equation 2.17.

To determine the fine structure of Δ we first restrict ourselves to the subspace $\mathcal{H}_{ij} = \mathcal{H}_{ij}^- \oplus \mathcal{H}_{ij}^+$ so that

$$M_{ij} := \Delta|_{\mathcal{H}_{ij}^-} : \mathcal{H}_{ij}^- \longrightarrow \mathcal{H}_{ij}^+. \tag{2.19}$$

Without loss of generality we choose the representation restricted to $\mathcal{H}_{ij}$ to be

$$\rho(a_i, a_j) = a_i \otimes 1_{n_j} \oplus 1_{n_i} \otimes a_j, \tag{2.20}$$

which corresponds to the choice of equation (2.7) with multiplicity $1_{m_{ij}} = 1$. For the restricted Dirac operator, real structure and chirality we choose the following:

$$\mathcal{D}_{ij} = \begin{pmatrix} 0 & M_{ij} \\ \overline{M_{ij}} & 0 \end{pmatrix}, \quad J_{ij} = \begin{pmatrix} 0 & 1_{n_i+n_j} \\ 1_{n_i+n_j} & 0 \end{pmatrix} \circ \mathrm{cc}, \quad \chi_{ij} = \begin{pmatrix} (-1)1_{n_i+n_j} & 0 \\ 0 & 1_{n_i+n_j} \end{pmatrix}. \tag{2.21}$$

Keeping in mind that we are for reasons of simplicity dealing with real and quaternionic matrix algebras, the first order axiom in the form of equation 2.17 reads

$$M_{ij}\,(b_i \otimes a_j) - (a_i \otimes 1_{n_j})\,M_{ij}\,(b_i \otimes 1_{n_j}) - (1_{n_i} \otimes b_j)\,M_{ij}\,(1_{n_i} \otimes a_j) + (a_i \otimes b_j)\,M_{ij} = 0 \tag{2.22}$$

for all $a_i, b_i \in \mathcal{A}_i$ and $a_j, b_j \in \mathcal{A}_j$ with $i \neq j$. In particular we can therefore choose $a_i = b_i = 0$ and $a_j = b_j = 1_{n_j}$ which leads to $M_{ij} = 0$.

As a first result from this analysis we conclude therefore that the Dirac operator cannot map subspace to each other which are also mapped to each other by the real structure. This is a direct consequence of the orientability axiom which requires $i \neq j$ in 2.22. The same argument holds for the complex case and for any multiplicity m_{ij} in the representation 2.6.

Let us now consider the more general case where we restrict ourselves to the subspace

$$\tilde{\mathcal{H}} = \mathcal{H}_{ij}^- \oplus \mathcal{H}_{rs}^- \oplus \mathcal{H}_{ij}^+ \oplus \mathcal{H}_{rs}^+ \tag{2.23}$$

with the obvious restrictions of the real structure and the chirality. The sub-matrix Δ of the Dirac operator $\mathcal{D}$ restricted to $\tilde{\mathcal{H}}$ is

$$\Delta|_{\tilde{\mathcal{H}}} = \begin{pmatrix} 0 & M_{ij}^{rs} \\ M_{rs}^{ij} & 0 \end{pmatrix}, \tag{2.24}$$

with

$$M_{ij}^{rs} : \mathcal{H}_{ij}^- \to \mathcal{H}_{rs}^+ \quad \text{and} \quad M_{rs}^{ij} : \mathcal{H}_{rs}^- \to \mathcal{H}_{ij}^+. \tag{2.25}$$

From $\Delta = \Delta^t$ follows $(M_{ij}^{rs})^t = M_{rs}^{ij}$ and thus

$$\Delta|_{\tilde{\mathcal{H}}} = \begin{pmatrix} 0 & M_{ij}^{rs} \\ (M_{ij}^{rs})^t & 0 \end{pmatrix} = \begin{pmatrix} 0 & M \\ M^t & 0 \end{pmatrix}, \tag{2.26}$$

where we have dropped the indices to simplify the notation.

We have now to distinguish two classes of possible representations on $\tilde{\mathcal{H}}$. The first possible representation restricted to $\tilde{\mathcal{H}}$ is given by

$$\rho(a)|_{\tilde{\mathcal{H}}} = \rho^-(a_i, a_j) \oplus \rho^-(a_r, a_s) \oplus \rho^+(a_i, a_j) \oplus \rho^+(a_r, a_s) \tag{2.27}$$

with

$$\begin{aligned} \rho^-(a_i,a_j) &= a_i \otimes 1_{n_j}, \quad \rho^+(a_i,a_j) = 1_{n_i} \otimes a_j, \\ \rho^-(a_r,a_s) &= a_r \otimes 1_{n_s}, \quad \rho^+(a_r,a_s) = 1_{n_r} \otimes a_s, \end{aligned} \tag{2.28}$$

with $i \neq j$ and $r \neq s$. Plugging the representation and Δ into the first order condition 2.17 we find for the sub-matrix M the following equation

$$M\,(b_r \otimes a_s) - (a_i \otimes 1_{n_j})\,M\,(b_r \otimes 1_{n_s}) - (1_{n_i} \otimes b_j)\,M\,(1_{n_r} \otimes a_s) + (a_i \otimes b_j)\,M = 0. \tag{2.29}$$

This equation has two solutions with $M \neq 0$: $i = s$ with $\mathcal{A}_i = \mathcal{A}_j = \mathbb{R}, \mathbb{C}$ and $r = j$ with $\mathcal{A}_r = \mathcal{A}_j = \mathbb{R}, \mathbb{C}$. The possibilities i, j, r, s pair-wise different and $i = r$, $j = s$ lead to $M = 0$.

We have therefore the two possible representations with $M \neq 0$:

$$\begin{aligned} \rho^-(a_i,a_j) &= a_i 1_{n_j}, \quad \rho^+(a_i,a_j) = a_j, \\ \rho^-(a_r,a_i) &= a_r, \quad \rho^+(a_r,a_i) = a_i 1_{n_r}, \end{aligned} \tag{2.30}$$

for $i = s$ and

$$\begin{aligned} \rho^-(a_i,a_j) &= a_i, \quad \rho^+(a_i,a_j) = a_j 1_{n_i}, \\ \rho^-(a_j,a_s) &= a_j 1_{n_s}, \quad \rho^+(a_j,a_s) = a_s, \end{aligned} \tag{2.31}$$

for $r = j$. These two possibilities are obviously equivalent.

The second general class of representations is of the form

$$\begin{aligned} \rho^-(a_i,a_j) &= a_i \otimes 1_{n_j}, \quad \rho^+(a_i,a_j) = 1_{n_i} \otimes a_j, \\ \rho^-(a_r,a_s) &= 1_{n_r} \otimes a_s, \quad \rho^+(a_r,a_s) = a_r \otimes 1_{n_s}. \end{aligned} \tag{2.32}$$

Using again the first order axiom 2.17 we find

$$M\,(a_r \otimes b_s) - (a_i \otimes 1_{n_j})\,M\,(1_{n_r} \otimes b_s) - (1_{n_i} \otimes b_j)\,M\,(a_r \otimes 1_{n_s}) + (a_i \otimes b_j)\,M = 0. \tag{2.33}$$

For $i = r$ ($\mathcal{A}_i = \mathcal{A}_j$) this has a general solution of the form $M = 1_{n_i} \otimes m$ with $m \in M_{n_j \times n_s}(\mathbb{C})$ and for $j = s$ ($\mathcal{A}_j = \mathcal{A}_s$) the solution has the form $M = m \otimes 1_{n_j}$ with $m \in M_{n_i \times n_r}(\mathbb{C})$. One has to keep in mind that $i \neq j$ and $r \neq s$ are required from the orientability axiom. All other possible equalities for i, j, r, s lead again to $M = 0$.

We have therefore again two possible representations with $M \neq 0$:

$$\begin{aligned} \rho^-(a_i,a_j) &= a_i \otimes 1_{n_j}, \quad \rho^+(a_i,a_j) = 1_{n_i} \otimes a_j, \\ \rho^-(a_i,a_s) &= 1_{n_i} \otimes a_s, \quad \rho^+(a_i,a_s) = a_i \otimes 1_{n_s} \end{aligned} \tag{2.34}$$

for $i = r$ and

$$\begin{aligned} \rho^-(a_i,a_j) &= a_i \otimes 1_{n_j}, \quad \rho^+(a_i,a_j) = 1_{n_i} \otimes a_j, \\ \rho^-(a_r,a_j) &= 1_{n_r} \otimes a_j, \quad \rho^+(a_r,a_j) = a_r \otimes 1_{n_j}. \end{aligned} \tag{2.35}$$

for $j = s$. These considerations can be easily adapted to the complex case which leads to some extra conditions which are treated in detail in [5]. Further restrictions also emerge from the axiom of Poincaré duality. This will be treated in the section on Krajewski diagrams.

The restriction to the subspace $\tilde{\mathcal{H}}$ it is now straightforward to choose particle- and antiparticle-subspaces $\mathcal{H}^P$ and $\mathcal{H}^A$. We define

$$\mathcal{H}^P := \mathcal{H}^-_{ij} \oplus \mathcal{H}^+_{rs} \quad \text{and} \quad \mathcal{H}^A := \mathcal{H}^+_{ij} \oplus \mathcal{H}^-_{rs}, \tag{2.36}$$

and identify $\mathcal{H}^{-(+)}_{ij}$ with the left-handed (anti)particles and $\mathcal{H}^{+(-)}_{rs}$ with the right-handed (anti)particles. In this basis the Dirac operator takes the well known form

$$\mathcal{D} = \begin{pmatrix} \Delta^P & 0 \\ 0 & \overline{\Delta^P} \end{pmatrix} \quad \text{with} \quad \Delta^P = \begin{pmatrix} 0 & M \\ M^* & 0 \end{pmatrix}. \tag{2.37}$$

This split extends naturally to the whole Hilbert space but it is off course ambiguous to choose which subspace constitutes the particle space and which the antiparticle space. The main point is that this split is always possible which was not the case in KO-dimension zero without an S^0-real structure. The crucial difference to the case of KO-dimension zero with S^0-real structure is that in principle lepto-quark like mass terms may appear in the Dirac operator. These were forbidden by the S^0-real structure. The ambiguity in the choice of particle and antiparticles existed also in KO-dimension zero due to a freedom of choice for the S^0-real structure.

Finally we would like to give the general form of the chirality after splitting the Hilbert space into particle- and antiparticle-subspaces, $\mathcal{H} = \mathcal{H}^P \oplus \mathcal{H}^A$:

$$\chi \;=\; (\oplus_{i,j=1}^N 1_{(n_i)} \otimes \chi_{ji} 1_{m_{ji}} \otimes 1_{(n_j)}) \;\oplus\; (\oplus_{i,j=1}^N 1_{(n_i)} \otimes (-\chi_{ji}) 1_{m_{ji}} \otimes 1_{(n_j)}), \tag{2.38}$$

where $\chi_{ij} = \mp 1$ according to our previous convention on left-(right-)handed subspaces. The real structure is invariant under this change of basis.

Comparing to the classification of finite spectral triples [11, 5] we find ourselves in the position to adapt directly the arguments which lead to the diagrammatic approach of Krajewski. For the choice of particle- and antiparticle subspaces we will use the conventions of [1].

3 Irreducibility and Krajewski diagrams

We are again dealing with irreducible spectral triples so let us recall the basic definitions.

3.1 Irreducibility

Definition 3.1. i) A spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is *degenerate* if the kernel of $\mathcal{D}$ contains a non-trivial subspace of the complex Hilbert space $\mathcal{H}$ invariant under the representation ρ on $\mathcal{H}$ of the real algebra $\mathcal{A}$.
ii) A non-degenerate spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is *reducible* if there is a proper subspace

$\mathcal{H}_0 \subset \mathcal{H}$ invariant under the algebra $\rho(\mathcal{A})$ such that $(\mathcal{A}, \mathcal{H}_0, \mathcal{D}|_{\mathcal{H}_0})$ is a non-degenerate spectral triple. If the triple is real and even, we require the subspace $\mathcal{H}_0$ to be also invariant under the real structure J and under the chirality χ such that the triple $(\mathcal{A}, \mathcal{H}_0, \mathcal{D}|_{\mathcal{H}_0})$ is again real and even.

Krajewski and Paschke & Sitarz have classified all finite, real spectral triples [11, 5]. Let us summarize the basics of this classification using Krajewski's diagrammatic language.

3.2 Conventions and multiplicity matrices

We will again only treat the easy case, $\mathbb{K} = \mathbb{R}, \mathbb{H}$ in all components of the algebra to emphasise the differences with respect to the case of KO-dimension zero. For further details on the complex case and on multiple arrows we refer to [1].

We define the *multiplicity matrix* $\mu \in M_N(\mathbb{Z})$, N being the number of summands in $\mathcal{A}$, such that $\mu_{ij} := \chi_{ij}\, m_{ij}$, with m_{ij} being the multiplicities of the representation (2.6) and χ_{ij} the signs of the chirality (2.38). There are N minimal projectors in $\mathcal{A}$, each of the form $p_i = 0 \oplus \cdots \oplus 0 \oplus \mathrm{diag}(1_{(1)}, 0, ..., 0) \oplus 0 \oplus \cdots \oplus 0$. With respect to the basis p_i, the matrix of the intersection form is $\cap = \mu - \mu^T$, the relative minus sign has again its origin in the anti-commutation relation of the real structure J and the chirality χ.

If both entries μ_{ij} and μ_{ji} of the multiplicity matrix are non-zero, then they must have the opposite sign. This has again to be contrasted with the case in KO-dimension zero, where the same sign is required.

• Poincaré duality: The last condition to be satisfied by the multiplicity matrix reflects the Poincaré duality and requires the multiplicity matrix to obey $\det(\mu - \mu^T) \neq 0$. Since the intersection form is an anti-symmetric matrix, this readily restricts us to finite spectral triples with an even number of summands in the matrix algebra.

• The Dirac operator: The components of the (internal) Dirac operator are represented by horizontal or vertical lines connecting two nonvanishing entries of opposite signs in the multiplicity matrix μ and we will orient them from plus to minus. Each arrow represents a nonvanishing, complex submatrix in the Dirac operator: For instance μ_{ij} can be linked to μ_{ik} by



and this arrow represents respectively submatrices of M in $\mathcal{D}$ of type $m \otimes 1_{(n_i)}$ with m a complex $(n_j) \times (n_k)$ matrix.

Every arrow comes with three algebras: Two algebras that localize its end points, let us call them *right and left algebras* and a third algebra that localizes the arrow, let us call it *colour algebra*. For the arrow presented above the left algebra is $\mathcal{A}_j$, the right algebra is $\mathcal{A}_k$ and the colour algebra is $\mathcal{A}_i$.

We deduced however that if $i = j$ or $k = j$ the corresponding spectral triple does not satisfy the axiom of orientability, so the colour algebra must not coincide with the left of the right algebra. Translated into the language of Krajewski diagrams this means that the arrow must not touch the diagonal of the diagram.

The requirement of non-degeneracy of a spectral triple means that every nonvanishing entry in the multiplicity matrix μ is touched by at least one arrow. We will also restrict ourselves to minimal Krajewski diagrams. A minimal Krajewski diagram is defined in detail in [13], in short it means that it is not possible to remove an arrow from the diagram without changing the multiplicity matrix.

• Convention for the diagrams: Our arrows always point from right chirality for particles and antiparticles, to left chirality for particles and antiparticles. As a further convention the horizontal arrows will encode particles and the vertical arrows encode antiparticles. This choice is of course arbitrary. As in the case of the classification of finite spectral triples of KO-dimension zero [1, 2, 3, 4] there may appear "corners", i.e. a horizontal arrow and a vertical arrow connected to a single point. But since every arrow comes with its transposed arrow (through the transposed multiplicity matrix), we can choose here as well one pair of arrows to represent the particles and the other to represent the antiparticles.

4 The Classification

As mentioned in the introduction, we will not give a complete derivation of the physical content for the irreducible, minimal Krajewski diagrams under consideration. For this we refer to [4], where all the details of the resulting physical models can be found. As we will see, the Krajewski diagrams of the case with KO-dimension six form a subset of the diagrams found in KO-dimension zero. Since the diagrams are taken to correspond to irreducible spectral triples, only the first fermion family is contained as a physical model. Further families have to be added by hand, these spectral triples are no longer irreducible.

The minimal diagrams for the case of four summands in the matrix algebra were computed with a computer program based on the algorithm presented in [13]. Only the calculation for the intersection form was changed so that the condition $\det(\mu - \mu^T) = 0$ has to hold for the multiplicity matrix corresponding to the Krajewski diagram. Thus also diagrams with arrows touching the diagonal appear in the output of the algorithm and have to be removed by hand since they do not satisfy the axiom of orientability.

In the classification in [1, 2, 3, 4] all models were discarded that have either

- a dynamically degenerate fermionic mass spectrum,
- Yang-Mills or gravitational anomalies,
- a fermion multiplet whose representation under the little group is real or pseudo-real,
- or a massless fermion transforming non-trivially under the little group.

Checking the Krajewski diagrams whether or not they meet the above conditions is completely analogous to the KO-dimension zero case. In fact the change of KO-dimension does not affect the representation, the Dirac operator or the physical models produced by a diagram. The only item of the spectral triple which is changed is the chirality.

The multiplicity matrix is anti-symmetric so the Poincaré duality can only be satisfied if the number of summands in the matrix algebra is even. The classification will be done for the cases with two summands and four summands. A classification beyond four summands is currently in progress.

4.1 Two Summands

In the case of two summands only one minimal Krajewski diagram exists:



Since the arrow touches the diagonal the diagram cannot represent a spectral triple which obeys to the orientability axiom. Therefore it will be discarded.

4.2 Four Summands

We will now analyse the diagrams produced by the computer program. They form a subset of the Krajewski diagrams obtained for the four-summand case in KO-dim. zero treated in [4]. This result is far from obvious, for one might expect that new diagrams are possible due to the sign change in the multiplicity matrix. As mentioned above the physical models obtained from a given diagram do not depend on the KO-dimension. We will therefore just give the number of the corresponding diagram in [4] to simplify the comparison for the reader. Furthermore we will quickly summarise the main results from [4] for each diagram and give the reason why they fail to meet the classification requirements or, if they survive which physical models they produce.

Diagram 1:



This diagram corresponds to diagram 5 in [4]. It fails for it has no unbroken colour and is dynamically degenerate.

Diagram 2:



It corresponds to diagram 6 in [4] and fails because it has no unbroken colours and all the summands in the matrix algebra have to be 1-dimensional.

Diagram 3, 4:





They correspond to diagram 8 in [4]. It has as its corresponding model the electro-strong model which is treated in detail in [4].

Diagram 5:



This diagram corresponds to diagram 20 in [4]. It fails for it either exhibits a trivial little group or a charged neutrino.

Diagram 6, 7:



These models correspond to the diagrams 18 and 19 in [4]. They reproduce the standard model of particle physics with various possibilities for the colour groups and certain sub-models. For the standard model algebra we find

$$\mathcal{A}_{SM} = \mathbb{C} \oplus \mathbb{H} \oplus M_C(\mathbb{C}) \oplus \mathbb{C}, \tag{4.1}$$

where C is the number of colours which has to be fixed by hand. All the physical models produced by this diagram are treated in great detail in [4]. It is interesting to note that the maximal Dirac operator corresponding to the left diagram (diagram 18 in [4]) has the Standard Model Dirac operator as its *maximal* Dirac operator. It does not contain any Lepto-Quarks!

5 Conclusions

This classification shows that the standard model takes an even more prominent place among the finite spectral triples when passing from KO-dimension zero to KO-dimension six, cutting down the number of relevant Krajewski diagrams from 66 to seven. It is most interesting to note that the Krajewski diagrams in KO-dimension six form a proper sub-set of the Krajewski diagrams obtained in [4] for KO-dimension zero. We could not show whether this is a general feature or coincidence.

Although one has to content oneself with a minimal version of the standard model, not allowing for massive neutrinos in all generations and prohibiting Majorana-masses for the right-handed neutrino and thus the see-saw-mechanism, this is still consistent with experimental data. Furthermore the fermion-doubling problem is resolved, as was shown in [6, 7].

An important result is that a real, finite spectral triple in KO-dimension six allows a natural, though ambiguous split of the Hilbert space into left-(right-)handed particle and antiparticle subspaces. One can therefore reduce the input for the spectral triple by the S^0-real structure which was needed in the case of KO-dimension zero. This reduction of input results in a remarkable reduction of the number of possible Krajewski diagrams and consequently a reduction of the number of possible physical models.

Extending the standard model by introducing massive right-handed neutrinos, as done by Alain Connes [6] and John Barrett [7], necessitates in a modification of the axioms of noncommutative geometry, especially the orientability axiom [10].

Acknowledgements: The authors would like to thank T. Schücker for careful proof reading, Bruno Iochum and Thomas Krajewski for helpful discussions and Alain Connes for insightful comments. We gratefully acknowledge a scholarship of the Friedrich Ebert-Stiftung for J.-H. Jureit and a fellowship of the Alexander von Humboldt-Stiftung for C.A. Stephan.

References

[1] B. Iochum, T. Schücker, and C. A. Stephan. On a classification of irreducible almost commutative geometries. *J. Math. Phys.*, 45:5003–5041, 2004.

[2] J.-H. Jureit and C. A. Stephan. On a classification of irreducible almost commutative geometries, a second helping. *J. Math. Phys.*, 46:043512, 2005.

[3] T. Schücker. Krajewski diagrams and spin lifts. 2005.

[4] J.-H. Jureit, T. Schücker, and C. A. Stephan. On a classification of irreducible almost commutative geometries iii. *J. Math. Phys.*, 46:072303, 2005.

[5] T. Krajewski. Classification of finite spectral triples. *J. Geom. Phys.*, 28:1–30, 1998.

[6] A. Connes. Noncommutative geometry and the standard model with neutrino mixing. 2006.

[7] J. W. Barrett. A lorentzian version of the non-commutative geometry of the standard model of particle physics. 2006.

[8] A. Connes. *Noncommutative geometry.* Academic Press, London and San Diego, 1994.

[9] A. Connes. Gravity coupled with matter and the foundation of non-commutative geometry. *Commun. Math. Phys.*, 182:155–176, 1996.

[10] A. Connes. Private communication.

[11] M. Paschke and A. Sitarz. Discrete sprectral triples and their symmetries. *J. Math. Phys.*, 39:6191–6205, 1998.

[12] A. Connes. Noncommutative geometry and reality. *J. Math. Phys.*, 36:6194–6231, 1995.

[13] J.-H. Jureit and C. A. Stephan. Finding the standard model of particle physics: A combinatorial problem. *J. Comp. Phys.*, accepted for publication, 2005.

[14] C.A. Stephan. Almost-commutative geometry, massive neutrinos and the orientability axiom in KO-dimension 6. hep-th/0610097, 2006